

**Time limit:** 50 minutes.

**Instructions:** This test contains 10 short answer questions. All answers must be expressed in simplest form unless specified otherwise. Only answers written on the answer sheet will be considered for grading.

**No calculators.**

- Alice and Bob are painting a house. If Alice and Bob do not take any breaks, they will finish painting the house in 20 hours. If, however, Bob stops painting once the house is half-finished, then the house takes 30 hours to finish. Given that Alice and Bob paint at a constant rate, compute how many hours it will take for Bob to paint the entire house if he does it by himself.
- Compute  $9^6 + 6 \cdot 9^5 + 15 \cdot 9^4 + 20 \cdot 9^3 + 15 \cdot 9^2 + 6 \cdot 9$ .
- Let  $x_1$  and  $x_2$  be the roots of  $x^2 - x - 2014$ , with  $x_1 < x_2$ . Let  $x_3$  and  $x_4$  be the roots of  $x^2 - 2x - 2014$ , with  $x_3 < x_4$ . Compute  $(x_4 - x_2) + (x_3 - x_1)$ .
- For any 4-tuple  $(a_1, a_2, a_3, a_4)$  where each entry is either 0 or 1, call it *quadratically satisfiable* if there exist real numbers  $x_1, \dots, x_4$  such that  $x_1x_2^2 + x_2x_4 + x_3 = 0$  and for each  $i = 1, \dots, 4$ ,  $x_i$  is positive if  $a_i = 1$  and negative if  $a_i = 0$ . Find the number of *quadratically satisfiable* 4-tuples.
- $a$  and  $b$  are nonnegative real numbers such that  $\sin(ax + b) = \sin(29x)$  for all integers  $x$ . Find the smallest possible value of  $a$ .

- Find the minimum value of

$$\frac{1}{x-y} + \frac{1}{y-z} + \frac{1}{x-z}$$

for reals  $x > y > z$  given  $(x-y)(y-z)(x-z) = 17$ .

- Compute the smallest value  $p$  such that, for all  $q > p$ , the polynomial  $x^3 + x^2 + qx + 9$  has exactly one real root.
- $P(x)$  and  $Q(x)$  are two polynomials such that

$$P(P(x)) = P(x)^{16} + x^{48} + Q(x).$$

Find the smallest possible degree of  $Q$ .

- Let  $b_n$  be defined by the formula

$$b_n = \sqrt[3]{-1 + a_1 \sqrt[3]{-1 + a_2 \sqrt[3]{-1 + \dots a_{n-1} \sqrt[3]{-1 + a_n}}}}$$

where  $a_n = n^2 + 3n + 3$ . Find the smallest real number  $L$  such that  $b_n < L$  for all  $n$ .

- Let  $x_0 = 1, x_1 = 0$ , and  $x_i = -3x_{i-1} + x_{i-2}$  for  $i \geq 2$ . Let  $y_0 = 0, y_1 = 1$ , and  $y_i = -3y_{i-1} + y_{i-2}$  for  $i \geq 2$ . Compute

$$\sum_{i=0}^{2013} \frac{(x_i y_{2014} - y_i x_{2014})^2}{y_{2014}^2}.$$

You may give your answer in terms of at most ten values of the  $x_i$  and/or  $y_i$  (but must otherwise simplify completely).